

## Connectivity

### 5.1 Cut Vertices

**Definition 1.** A vertex  $v$  in a connected graph  $G$  is a *cut vertex* if  $G - v$  is disconnected.

**Definition 2.** A vertex  $v$  in a graph  $G$  is a *cut vertex* if  $G - v$  has more connected components than  $G$ .

**Exercise.** Prove that these two definitions are equivalent.

**Lemma** (End-Vertex Lemma). *If  $v$  is an end vertex in a graph  $G$ , then  $v$  is not a cut vertex of  $G$ .*

*Proof.* Let  $v$  belong to the connected component  $H$  of  $G$ . Grow a maximal path  $P$  starting from  $v$ . Since  $\deg v = 1$ , vertex  $v$  is one of the two ends of  $P$ . By the Maximal Path Theorem,  $H - v$  is connected. Thus  $v$  is not a cut vertex.  $\square$

**Theorem** (CZ, Theorem 5.1). *Let  $G$  be a graph containing a bridge  $e$  incident with vertex  $v$ . Vertex  $v$  is a cut vertex if and only if  $\deg v \geq 2$ .*

*Proof.* Let  $G$  be a graph containing a bridge  $e$  incident with vertex  $v$ .

$\Rightarrow$ : Suppose  $\deg v < 2$ . Since  $e$  is incident with  $v$ , we have  $\deg v \geq 1$ . Thus,  $\deg v = 1$ . By the End-Vertex Lemma,  $v$  is not a cut vertex.

$\Leftarrow$ : Suppose  $\deg v \geq 2$  but  $v$  is not a cut vertex. Let bridge  $e$  join  $v$  to  $w$  and let  $H$  be the connected component of  $G$  that contains  $v$ . Since  $\deg v \geq 2$ , vertex  $v$  is adjacent to some vertex  $x$  that is different from  $v$  or  $w$ . Therefore, in  $H$ , vertex  $w$  is connected to vertex  $x$  via the path  $P : w, v, x$ . This says that vertices  $v$ ,  $w$ , and  $x$  are all in the same component  $H$ . By assumption, vertex  $v$  is not a cut vertex of  $G$ ; thus  $v$  is not a cut vertex of  $H$ , i.e.,  $H - v$  is connected. This says that all vertices in  $H - v$  are in the same component. Hence, vertex  $w$  is connected in  $H - v$  to vertex  $x$  via some path  $Q$ . Concatenating  $P$  to  $Q$  gives a cycle in  $H$ . This cycle contains edge  $e$ , contradicting the fact that  $e$  is a bridge.  $\square$

**Corollary** (CZ, Corollary 5.2). *Let  $G$  be a connected graph of order 3 or more. If  $G$  contains a bridge, then  $G$  contains a cut-vertex.*

*Proof.* Let  $G$  be a connected graph of order at least 3 and let  $e = vw$  be a bridge in  $G$ . By the Bridge Lemma,  $G - e$  consists of two components: component  $G_v$  containing  $v$  and component  $G_w$  containing  $w$ . Since  $G$  has at least 3 vertices, at least one of  $G_v$  and  $G_w$  has more than one vertex. Assume wlog that  $G_v$  has more than one vertex. Thus,  $\deg_{G_v} v \geq 1$ . Since  $v$  is adjacent in  $G$  to  $w$  but  $w$  is not in  $G_v$ , we conclude that  $\deg_G v \geq 2$ . By Theorem 5.1  $v$  is a cut vertex (in  $G$ ).  $\square$

**Theorem** (CZ, Corollary 5.4). *A vertex  $v$  of a connected graph  $G$  is a cut vertex of  $G$  if and only if there exist vertices  $u$  and  $w$  distinct from  $v$  such that there's at least one  $u - w$  path in  $G$  and  $v$  lies on every  $u - w$  path in  $G$ .*

*Proof.* Suppose  $v$  is a cut vertex of a connected graph  $G$ . Then  $G - v$  is disconnected, i.e., it has at least 2 connected components. Let  $u$  be any vertex in  $G - v$  and let  $w$  be any other vertex in  $G - v$  belonging to some component different from  $u$ 's. Since  $G$  is connected, there exist some  $u - w$  path in  $G$ . However, there exist no  $u - w$  path in  $G - v$  because  $u$  and  $w$  come from different components of  $G - v$ . This implies that each  $u - w$  path in  $G$  passes through  $v$ , since  $G - v$  differs from  $G$  only in that  $G - v$  misses vertex  $v$  and all edges incident to  $v$ .

Conversely, suppose  $v$  is any vertex in a connected graph  $G$  and  $G$  contains some vertices  $u, w$  such that there's at least one  $u - w$  path in  $G$  and  $v$  lies on every  $u - w$  path in  $G$ . This assumption implies that any  $u - w$  path in  $G$  can no longer be a  $u - w$  path when vertex  $v$  and all its incident edges are deleted from  $G$ . Thus,  $G - v$  is disconnected since it has vertices  $u$  and  $w$  that are not connected. Hence,  $v$  is a cut vertex of  $G$ .  $\square$

**Theorem** (CZ, Corollary 5.6). *Every nontrivial connected graph contains at least two vertices that are not cut vertices.*

*Proof.* This is just Theorem 1.9 (with the phrase “connected graph of order 3 or more” changed to “connected nontrivial graph”) restated in terms of cut vertices. See Handout #5.  $\square$

## 5.2 Blocks

**Definition.** A *nonseparable graph* is nontrivial, connected, and has no cut vertex.

*Note:*  $K_2$  is the only nonseparable graph of order less than 3.

**Theorem** (CZ, Theorem 5.7). *A graph of order at least 3 is nonseparable if and only if every two vertices lie on a common cycle.*

*Proof.* Let  $G$  be a graph of order at least 3.

Suppose every two vertices of  $G$  lie on a common cycle. Graph  $G$  is nontrivial since its order is at least 3. Let  $x, y$  be two vertices of  $G$ . By assumption there is a cycle  $C$  that contains both  $x$  and  $y$ . Thus there is an  $x - y$  path along  $C$ . Hence  $G$  is connected. Fix a vertex  $v$ . Let  $u, w$  be any two vertices distinct from  $v$ . By assumption there is a cycle that contains both  $u$  and  $w$ . This cycle gives two internally disjoint  $u - w$  paths, at least one of which does not go through  $v$ . Hence,  $v$  is not a cut vertex by Corollary 5.4. Thus, graph  $G$  contains no cut vertex. Therefore,  $G$  is nonseparable.

Conversely, suppose  $G$  is nonseparable. Let  $u$  be a vertex of  $G$ . We will prove that if  $v$  is any vertex of  $G$  distinct from  $u$ , then there is a cycle that goes through both  $u$  and  $v$ , by induction on the distance  $d(u, v)$  between  $u$  and  $v$ . First suppose that  $d(u, v) = 1$ , i.e.,  $G$  has an edge  $e$  joining  $u$  to  $v$ . Since  $G$  has order at least 3 and it contains no cut vertex (because  $G$  is nonseparable), we conclude by Corollary 5.2 that  $G$  contains no bridge. Therefore,  $e$  is not a bridge; and thus some cycle  $C$  contains  $e$ . Hence, cycle  $C$  contains both  $u$  and  $v$  (since  $e$  joins  $u$  to  $v$ ). Next suppose that  $d(u, v) = k > 1$  and assume inductively that for any vertex  $x$ , where  $0 < d(u, x) < k$ , there exists some cycle that goes through both  $u$  and  $x$ . Let  $P : u = v_0, v_1, \dots, v_{k-1}, v_k = v$  be a  $u - v$  geodesic. Since  $0 < d(u, v_{k-1}) = k - 1 < k$ , there exists, by inductive assumption, a cycle  $C$  that passes through both  $u$  and  $v_{k-1}$ . If cycle  $C$  goes through  $v_k$  as well, then we are done. So assume from now on that  $C$  does not go through  $v_k$ . Since  $G$  contains no cut vertex,  $v_{k-1}$  is not a cut vertex. This means that there exists some path in  $G - v_{k-1}$  (and in  $G$  as well) connecting  $v_k$  to some vertex  $x \in V(C) \setminus v_{k-1}$ . Let  $Q$  be such a path of shortest possible length. Appending the  $x - v_{k-1}$  path in  $C$  that goes through  $u$  (this path is unique if  $x \neq u$ ) to  $Q$  gives a  $v - v_{k-1}$  path  $P'$  in  $G$  that goes through  $u$ . Path  $P'$  together with edge  $v_{k-1}v_k$  gives a cycle in  $G$  that goes through both  $u$  and  $v$ . Our claim follows by induction.  $\square$

**Definition.** Let  $G$  be a graph of positive size. Define a relation  $R$  on  $E(G)$  as follows. For any edges  $e$  and  $f$ , declare  $eRf$  iff  $e = f$  or there is a cycle in  $G$  that contains both  $e$  and  $f$ .

**Theorem** (CZ, Theorem 5.8). *The relation  $R$  is an equivalence relation.*

*Proof.* ... □

**Definition 1.** A *block* of  $G$  is a nonseparable subgraph of  $G$  that is not a proper subgraph of any other nonseparable subgraph of  $G$ .

**Definition 2.** A *block* of  $G$  is a subgraph of  $G$  induced by the edges in an equivalence class defined by the relation  $R$  defined above.

**Theorem** (CZ, Exercise 5.15). *The two definitions of block are equivalent.*

*Proof.* ... □

**Corollary** (CZ, Corollary 5.9). *Every two distinct blocks  $B_1$  and  $B_2$  in a nontrivial connected graph  $G$  have the following properties:*

- (a) *The blocks  $B_1$  and  $B_2$  are edge-disjoint.*
- (b) *The blocks  $B_1$  and  $B_2$  have at most one vertex in common.*
- (c) *If  $B_1$  and  $B_2$  have a vertex  $v$  in common, then  $v$  is a cut vertex of  $G$ .*

*Proof.* (a) By definition 2 of block, the edges  $E(B_1)$  of block  $B_1$  and the edges  $E(B_2)$  of block  $B_2$  belong to different equivalence classes. Therefore,  $E(B_1) \cap E(B_2) = \emptyset$ .

(b) Assume for the sake of contradiction that  $|V(B_1) \cap V(B_2)| \geq 2$ . Since  $B_1$  is connected (because it's a block), for any two vertices shared by the two blocks there exists a path in  $B_1$  connecting them. Let  $P_1$  be a shortest path in  $B_1$  connecting any two shared vertices; say that  $P_1$  connects  $v$  to  $w$ . Path  $P_1$  is nontrivial since  $v \neq w$ . Since  $B_2$  is connected (because it's a block), there exists a path  $P_2$  in  $B_2$  connecting  $v$  to  $w$ . Path  $P_2$  is nontrivial since  $v \neq w$ . Concatenating  $P_1$  to  $P_2$  gives a cycle containing some edge  $e_1$  in  $B_1$  and some edge  $e_2$  in  $B_2$ . Thus,  $e_1$  and  $e_2$  belong to the same block by definition 2 of block. This contradicts part (a).

(c) Let  $v \in V(B_1) \cap V(B_2)$ . Being a block,  $B_1$  is connected and nontrivial; thus, there exists  $u_1 \in V(B_1)$  adjacent to  $v$ . Being a block,  $B_2$  is connected and nontrivial; thus, there exists  $u_2 \in V(B_2)$  adjacent to  $v$ .

We'll show that every  $u_1 - u_2$  path in  $G$  contains  $v$ . Assume for the sake of contradiction that there is some  $u_1 - u_2$  path  $P$  in  $G$  not containing  $v$ . Path  $P$  together with  $v$  and edges  $u_1v$  and  $vu_2$  give a cycle  $C$  containing both  $u_1v$  and  $vu_2$ . Hence, edges  $u_1v$  and  $vu_2$  belong to the same equivalence class, i.e., same block. This contradicts part (a).

Therefore,  $v$  is a cut vertex of  $G$ .  $\square$

## 5.3 Connectivity

**Definitions.** Let  $G = (V, E)$  be a connected graph. A subset  $U$  of  $V$  is called a *vertex cut* if  $G - U$  is disconnected. A *minimum vertex cut* of  $G$  is a vertex cut of least cardinality. The (*vertex*) *connectivity*  $\kappa(G)$  of  $G$  is defined as follows: for a disconnected graph  $G$ ,  $\kappa(G) = 0$ ; for a connected graph  $G$ ,  $\kappa(G)$  equals the cardinality of a smallest vertex subset  $U$  such that  $G - U$  is either disconnected or trivial. A graph  $G$  is *k-connected* if  $\kappa(G) \geq k$ . Note that any graph  $G$  satisfies  $0 \leq \kappa(G) \leq n - 1$ . Note also that for a connected graph  $G$  of order  $n$ ,  $\kappa(G) = n - 1$  if and only if  $G \cong K_n$ .

Let  $G = (V, E)$  be a connected graph. A subset  $X$  of  $E$  is an *edge cut* if  $G - X$  is disconnected. An edge cut  $X$  is *minimal* if no proper subset of  $X$  is an edge cut. A *minimum edge cut* is an edge cut of minimum size.

Note that a minimal edge cut is not necessarily minimum, but every minimum edge cut is necessarily minimal. (Prove!)

The following lemma characterizes minimal edge cuts.

**Lemma** (Minimal Edge Cut Lemma). *If  $X$  is a minimal edge cut of a connected graph  $G$ , then  $G - X$  contains exactly 2 components. Moreover,  $X$  consists of all the edges of  $G$  that join a vertex in one component to a vertex in another component.*

*Proof.* Assume  $G$  is a connected graph and  $X$  is a minimal edge cut of  $G$ . Choose any edge  $e \in X$ , say  $e$  joins  $u$  to  $v$ . We have  $G - X$  is disconnected but  $G - (X \setminus \{e\})$  is connected. This means that  $e$  is a bridge in  $G - (X \setminus \{e\})$ . By the Bridge Lemma,  $G - (X \setminus \{e\}) - e$  consists of exactly two components, one containing  $u$  and the other containing  $v$ . Since  $G - (X \setminus \{e\}) - e = G - X$ , we conclude that  $G - X$  consists of exactly two components, say  $G_1$  and  $G_2$ , and every edge of  $X$  joins a vertex in  $G_1$  to a vertex in  $G_2$ .

Let  $e$  be any edge in  $G$  joining a vertex in  $G_1$  to a vertex in  $G_2$ . We'll show that  $e \in X$ . Suppose not. Then  $G - X$  contains edge  $e$ . This contradicts  $G_1$  and  $G_2$  are distinct components in  $G - X$ .  $\square$

The *edge connectivity*  $\lambda(G)$  of  $G$  is defined as follows: for a disconnected graph  $G$ ,  $\lambda(G) = 0$ ; for a connected graph  $G$ ,  $\lambda(G)$  equals the cardinality of a smallest edge subset  $X$  such that  $G - X$  is either disconnected or trivial. A graph  $G$  is  *$k$ -edge-connected* if  $\lambda(G) \geq k$ . Note that any graph  $G$  satisfies  $0 \leq \lambda(G) \leq n - 1$ .

**CZ, Example 5.10** Show that  $\lambda(K_n) = n - 1$ .

*Proof.* ...  $\square$

**Theorem** (CZ, Theorem 5.11). *For any graph  $G$ ,*

$$\kappa(G) \leq \lambda(G) \leq \delta(G).$$

*Proof.* ...  $\square$